

HIGH ELEVATIONS

High altitude illnesses can kill. Military operations occurring at elevations over 6,000 feet can seriously impact unit and individual effectiveness. Serious illness or death can result if you ascend rapidly without allowing for acclimatization. Remain well hydrated; individual water requirements are greater at higher altitudes.

- When deployed to high mountain areas, be observant of the common symptoms of mountain sickness: headache, nausea, vomiting, dizziness, fatigue, irritability, and coughing. Seek medical attention immediately if you experience any of these symptoms.
- Pyridostigmine bromide tablets may increase the chance of dizziness or fainting during the first 24 hours at high altitude if you are not acclimatized.
- Lower oxygen levels at high altitudes ("thin air") combined with the heavier work requirements when wearing mission-oriented protective posture (MOPP) gear can increase your risk of high altitude illnesses. When wearing MOPP gear at higher altitudes, you may require more time and concentration to perform assigned tasks.
- For appropriate countermeasures during high altitude operations, see GTA 08-05-062 and GTA 08-05-060, *A Soldier's Guide to Staying Healthy at High Elevations*.

HEARING PROTECTION

It is essential that you use properly fitted hearing protection during military operations. Exposure to high-intensity noise may cause hearing loss that can adversely affect your combat effectiveness and individual readiness. Good hearing is essential to mission success. If you are a dismounted soldier, the Combat Arms Earplug (NSN 6515-01-466-2710) will protect you from the impact noise of weapons fire while only slightly interfering with voice communications and detection of combat sounds such as vehicle noise, footfalls in leaves, and the closing of a rifle bolt. While not as effective as the Combat Arms Earplug in preserving your ability to hear important mission-related sounds, noise muffs or standard earplugs are very effective in preventing noise-induced injury. If you are a member of vehicle or helicopter crews, your combat vehicle crew or aircrew helmets have built-in hearing protectors.

ORAL HEALTH

Dental disease is a common problem during deployments due to the challenge of maintaining good oral hygiene. You should deploy with toothbrush, dental floss, and fluoride toothpaste. Daily flossing and twice daily brushing of teeth is the best way to ensure prevention of periodontal disease and to decrease your risk of problems such as trench mouth and tooth decay. In difficult tactical environments, teeth should be brushed at least once a day. Seek medical attention immediately at the onset of any dental problems.

PRE-DEPLOYMENT HEALTH INFORMATION

- Complete the Pre-Deployment Health Assessment (DD FORM 2795) to assess your state of health before deployment and to assist health care providers in identifying your medical needs and providing present and future medical care to you.
- You will not have access to your health care record during the deployment. The Adult Preventive and Chronic Care Flowsheet (DD FORM 2766) will be used as your deployment health record. This document will include information on all your immunizations, any medications you are currently taking, and any ongoing medical problems that you may have. When you go through readiness processing, ensure that all appropriate information is documented on your DD FORM 2766. When you return home, this information will be placed in your regular health record.

INFORMATION ABOUT YOUR HEALTH CARE WHILE DEPLOYED

- It is important that you know where to seek health care while deployed. This may or may not be through the same channels as your home station. Ask your chain of command for more information.
- While deployed, you must maintain your health and seek care whenever an illness or injury threatens your ability to complete your tasks. Your unit is depending on you. It is always better to seek care early so that your problems can be documented appropriately and taken care of immediately.

POST-DEPLOYMENT HEALTH INFORMATION

- Complete the Post-Deployment Health Assessment (DD FORM 2796) to assess your state of health after deployment and to assist health care providers in identifying your medical needs and providing present and future medical care to you.
- If you become sick after you return home, tell your physician that you were deployed.
- After deployment to Honduras, it is especially important to complete malaria medications as directed, and receive follow-on medical care/tests as directed.

Contact your Preventive Medicine or Medical Support Unit for more information.

DISTRIBUTION: UNLIMITED

Prepared by:



U.S. Army Center for Health Promotion & Preventive Medicine
<http://chppm-www.apgea.army.mil>

SIPRNet: <http://usachppm1.army.smil.mil>

(800) 222-9698/ DSN 584-4375/(410) 436-4375

SHG 009-0302

A SOLDIER'S GUIDE TO STAYING HEALTHY IN HONDURAS

This country-specific guide should be used in conjunction with GTA 08-05-062, *Guide to Staying Healthy*, and is intended to provide information that can help reduce your risk of Disease and Non-battle Injuries (DNBI) when deployed. This health threat and countermeasure information is based on the most current data available from U.S. Department of Defense medical agencies at the time of production. In addition to the information in this guide, you should also receive force health protection, health threat, and preventive medicine countermeasures training/briefings prior to and, as required, throughout the length of your deployment.

HONDURAS OVERVIEW

Honduras, slightly larger than Tennessee, is located in Central America and is one of the four poorest nations in the Western Hemisphere. Narrow coastal plains extend inland about 75 miles at the widest point. These lowlands consist of grasslands, swamps, pine forests, and tropical rain forests. The mountainous interior comprises nearly 80 percent of the country. The borders of the country include Nicaragua in the east, the Pacific Ocean and Nicaragua in the south, El Salvador and Guatemala in the west, and the Caribbean Sea in the north. The elevation ranges from sea level along the Caribbean Sea to over 9,000 feet in the southern volcanic highland area; however, the highland elevation generally is between 7,300 and 8,600 feet. The Caribbean Lowlands have a tropical wet climate with consistently high temperatures and humidity and rainfall evenly distributed throughout the year. The Pacific Lowlands have a tropical wet climate with high temperatures but a distinct dry season from November through April. The interior highlands also have a distinct dry season, but temperatures decrease as elevation increases. The northern coastal region receives an average annual rainfall of 70 to 100 inches while the Pacific coastal plains receive somewhat less. The inland areas generally receive a more moderate rainfall of 33 inches annually. The average temperature range is 61° to 81° F throughout the year. Above 6,500 feet, temperatures can fall to near freezing at night, and frost sometimes occurs. Earthquakes are frequent but generally mild. Damaging hurricanes and tropical storms with flooding are common and can occur from the summer through the fall.

HONDURAS RISK ASSESSMENT

Based on a combination of all major infectious diseases that occur in a country, an overall country risk level is assigned as low, intermediate, high, or highest risk. This region is INTERMEDIATE RISK for infectious diseases. Diseases of military importance to forces deployed to Honduras include hepatitis A and E, typhoid fever and several diarrheal

HONDURAS RISK ASSESSMENT (CONTINUED)

diseases to include cholera, all acquired by consuming contaminated food, water, and dairy products; vector-borne diseases such as malaria, dengue fever, Eastern equine encephalitis, Rocky Mountain spotted fever, American trypanosomiasis (Chagas disease), and leishmaniasis, which are acquired through the bites of insects and ticks; leptospirosis from swimming, wading, or skin contact with contaminated water; rabies and Q fever from animal contact; tuberculosis from contact with human respiratory secretions or droplets; and sexually transmitted diseases. Environmental factors also pose a significant health risk to deployed forces and include sewage, agricultural, and industrial contamination of water and food supplies and localized air pollution. Additionally, high altitude illnesses are a potential DNBI in the mountainous regions.

INCREASED REGIONAL DISEASE THREATS

Recent hurricane and tropical storm destruction of an already weak Honduras public infrastructure increases the potential for infectious disease transmission throughout the country. Losses from the most recent storm include 35 water systems and major access roads.

There is significant risk in this country for acquiring incapacitating mosquito-borne diseases including dengue fever and malaria. Both are transmitted by mosquitoes that exist throughout the country, year-round, in both urban and rural areas, and that feed day and night, indoors or out. The risk for dengue fever is highest in urban environments; the risk for malaria is highest in rural areas. Large epidemics of dengue fever occur at irregular intervals. Risk of dengue fever is increased during the rainy months when the mosquito population increases, particularly in northern coastal urban areas. In the past, malaria rates have increased tenfold or more following hurricanes in the region. The risk for malaria is highest in the northern coastal lowlands and along the border with Nicaragua. Preventing exposure to mosquitoes and other biting vectors at all times and in all areas will help reduce your risk of acquiring vector-borne diseases.

Increased potential for hookworm infection is present in the region. Hookworm infections are caused by a parasite in the soil that penetrates the skin, usually the foot, and causes skin irritation, anemia, and respiratory and intestinal symptoms. Hookworm infections are usually acquired by walking barefoot.

FIELD SANITATION TEAM

Each company-sized unit has a Field Sanitation Team (FST) whose members are trained (40-hour course) and fully equipped IAW AR 40-5, FM 4-25.12, and FORSCOM REG 700-2. Know who the members of your FST are, and know how they can assist in preventing medical threats to your health. Become familiar with FST equipment and training.

VECTOR-BORNE DISEASES

There are several vector-borne diseases present in Honduras. They include malaria, dengue fever, and Eastern equine encephalitis from mosquitoes; Rocky Mountain spotted fever from ticks; leishmaniasis from sand flies; and American trypanosomiasis from "kissing bugs." There are many other diseases spread by the bites of mosquitoes, ticks, sand flies, fleas, mites, and lice. Your local medical authority will determine if these diseases or other vector-borne diseases are a threat in your specific location and provide appropriate countermeasures.

- Take your malaria prevention pills when directed to do so. This is CRITICAL. Malaria is widespread in this region. Normally, you will begin taking medication prior to arriving in the area, while in the area, and after returning home.
- Use the DOD Insect Repellent System detailed in GTA 08-05-062 to reduce your risk of acquiring a vector-borne disease. Wear permethrin-treated uniforms with trousers tucked into boots, sleeves rolled down, and undershirt tucked into trousers. Wear DEET on exposed skin.
- When deployed to this region, sleep under a permethrin-treated bed net to repel insects and further reduce risks of vector-borne diseases. Many insects in this region feed during the night, including mosquitoes that transmit malaria.
- When using both DEET and sunscreen products, apply sunscreen to the skin first so it does not interfere with the effectiveness of the DEET. After 30 minutes to an hour, apply the DEET. This allows the sunscreen to penetrate and bind to the skin first.

FOOD-BORNE AND WATER-BORNE DISEASES

In the past 5 years, hurricanes have caused severe localized contamination of water systems, particularly in the northern areas. The diseases of greatest risk throughout the region, in both rural and urban areas, are bacterial diarrhea, hepatitis A, and typhoid fever associated with contaminated food, water, and/or ice. Do not consume any food, water, or beverages (to include bottled water) that have not been approved by the U.S. military. Assume all non-approved food, ice, and water is contaminated. Water and food items available in this region, including dairy products, fish, fruits, and vegetables, may contain unsafe levels of pesticides, chemical fertilizers, bacteria, and viruses. Contamination with human or animal waste is widespread. Even a one-time consumption of these foods or water may cause severe illness. See GTA 08-05-062 for appropriate countermeasures.

SEXUALLY TRANSMITTED DISEASES

Sexually transmitted diseases are highly prevalent in this region. Gonorrhea, chlamydia, and other infections are common, and may affect a high percentage of personnel who have sexual contact. Human immunodeficiency virus (HIV) and hepatitis B also occur. Though the immediate impact of HIV and hepatitis B on an operation is limited, the long-term impact on your individual health is substantial. See GTA 08-05-062 for appropriate countermeasures.

HOT AND COLD WEATHER INJURIES

Heat injury can occur in individuals not acclimated to the climate. Heat injuries are more likely to occur during the early phase of deployment; acclimatization is critical. Individual and unit countermeasures are extremely important. Cold injuries can also occur in the mountainous areas of this region. The effects of cold weather are more severe in high mountainous areas due to reduced oxygen and lower air pressure. When deploying to the mountainous areas in this region, check with your unit on the requirement for packing the extended cold weather clothing system. See GTA 08-05-062 for appropriate countermeasures.

HAZARDOUS ANIMALS AND PLANTS

- Several species of highly poisonous snakes, which are well camouflaged and very aggressive, live in the country. Consider any snake encountered as poisonous, and do not handle. Seek immediate medical attention if bitten; untreated snakebites may cause serious illness or death within 1 hour.
- Several species of spiders capable of inflicting a painful bite are present throughout the country (including some very large and physically imposing tarantulas). If possible, avoid sleeping on the ground. Shake out boots, bedding, and clothing prior to use, and never walk barefoot. If bitten or stung, seek medical attention immediately.
- Some regional plants have thorns, stinging hairs, or toxic resins that may puncture the skin or introduce poison into the skin causing skin irritation, rashes or infections. Avoid skin contact with plants when tactically feasible.
- Contact with the smoke from the burning of these plants can also cause skin rashes and damage to your lungs.
- Clean your clothing after contact with harmful plants. Decontaminate clothing by washing with soap and water.
- Some regional plants may cause systemic poisoning if leaves, berries, flowers, or other plant parts are chewed or swallowed. Symptoms include dizziness, vomiting, irregular heartbeat, and delirium or hallucinations.

TOXIC INDUSTRIAL CHEMICALS AND MATERIALS

When deployed, you may face health risks from industrial chemicals and materials as a result of activities by terrorists or warring parties; accidents related to improper design, maintenance, or operation of indigenous industrial facilities; inadvertent exposure to toxic waste materials in the environment; or improper handling or disposal of hazardous material with which our own forces deploy. The degree of health risks depends upon many parameters. Consult your medical authority for additional information.

SKIN DISEASES

Skin irritations and infections, such as athlete's foot and ringworm, are common medical threats during any deployment and are commonly caused by fungi. The best prevention is to maintain clean, dry skin. See GTA 08-05-062 for additional countermeasure information.